

USAGE

USAGE AREA	USAGE TIME	USAGE AND AMOUNT	
		BY LEAVES	BY SOIL
All greenhouse vegetable growing (Tomatoes, peppers, eggplants, cucumbers, strawberries, etc.)	It is applied with an interval of one week from planting to the end of harvest.	200-250 cc in 100 liters of water	1000-1500 cc/da
All open field vegetable growing (Tomatoes, peppers, eggplants, cucumbers, strawberries, etc.)	It is applied 2-3 times with an interval of 15-20 days from planting to the end of harvest.	250-300 cc in 100 liters of water	1000-1500 cc/da
All winter vegetables with edible leaves (cauliflower, leek, spinach, lettuce, curly lettuce, iceberg lettuce, etc.)	It is applied 2-3 times with an interval of 20-30 days from planting to the end of harvest	250-300 cc in 100 liters of water	1000-1500 cc/da
Cut Flowers Growing	It is applied weekly throughout the season beginning from planting.	200-250 cc in 100 liters of water	1000-1500 cc/da
Melons, Watermelons, Pumpkins, etc.	It is applied 2-3 times with an interval of 20-30 days beginning from planting.	250-300 cc in 100 liters of water	1000-1500 cc/da
All Tubers (Onions, Garlic, Potatoes, Carrots, Radish, Sugar Beet)	It is applied at the beginning of flowering and in the period of tuber binding and tuber growth.	200-250 cc in 100 liters of water	1000-1500 cc/da
All Fruit Trees (Apples, Pears, Peaches, Apricots, Quinces, Cherries, Almonds, Grapes, etc.)	At the beginning of blooming, fruit formation, fruit growing and before harvest.	250-300 cc in 100 liters of water	1000-1500 cc/da or 75-100 cc/tree
Bananas, Citrus, Figs, Olives, Walnuts, Almonds, Kiwis	At the beginning of blooming, fruit formation, fruit growing and before harvest.	250-300 cc in 100 liters of water	1000-1500 cc/da or 75-100 cc/tree
Industrial Crops and Legume (Sunflowers, Corns, Chickpeas, Soy, Beans, Peas, Peanuts, Lentil, Tea)	After the plants reach 10-15 cm in height, it is applied 2-3 times with an interval of 20 days.	200-250 cc in 100 liters of water	1000-1500 cc/da
Field Crops (Barley, Wheat, Rye, Paddy)	It is applied during tillering and bolting period.	200-250 cc in 100 liters of water	-----
Green Fields and Forage Plants (Clover, Sainfoin, Vetch)	It is applied 10 days after each mowing.	200-250 cc in 100 liters of water	-----