



BLACKKING

Blackking is a special product containing organic substance, organic carbon and amino acids that can be applied from the soil and leaves. It enhances the soil structure. It enables the growth of microorganisms in the soil and the transport of nutrients and water that cannot be taken by the plant.

It increases the photosynthesis of plants and supports the yield between 20% and 50%.

Due to its natural structure, it quickly carries macro and micro nutrients in the soil to the plant.

It supports root development. In foliar use, it provides homogeneous flower and fruit formation and earliness.

GUARANTEED SUBSTANCE	% W/W
Organic Matter	35
Organic Carbon	13
Organic Nitrogen	3
Water-soluble Potassium Oxide (K ₂ O)	2
Free Amino Acids	3
pH interval	4-6

USAGE

USAGE AREA	USAGE TIME	USAGE AND AMOUNT	
		BY LEAVES	BY SOIL
All greenhouse vegetable growing (Tomatoes, peppers, eggplants, cucumbers, strawberries, etc.)	It is applied with an interval of one week from planting to the end of harvest.	200-250 cc in 100 liters of water	1000-1500 cc/da
All open field vegetable growing (Tomatoes, peppers, eggplants, cucumbers, strawberries, etc.)	It is applied 2-3 times with an interval of 15-20 days from planting to the end of harvest.	250-300 cc in 100 liters of water	1000-1500 cc/da
All winter vegetables with edible leaves (cauliflower, leek, spinach, lettuce, curly lettuce, iceberg lettuce, etc.)	It is applied 2-3 times with an interval of 20-30 days from planting to the end of harvest	250-300 cc in 100 liters of water	1000-1500 cc/da
Cut Flowers Growing	It is applied weekly throughout the season beginning from planting.	200-250 cc in 100 liters of water	1000-1500 cc/da
Melons, Watermelons, Pumpkins, etc.	It is applied 2-3 times with an interval of 20-30 days beginning from planting.	250-300 cc in 100 liters of water	1000-1500 cc/da
All Tubers (Onions, Garlic, Potatoes, Carrots, Radish, Sugar Beet)	It is applied at the beginning of flowering and in the period of tuber binding and tuber growth.	200-250 cc in 100 liters of water	1000-1500 cc/da
All Fruit Trees (Apples, Pears, Peaches, Apricots, Quinces, Cherries, Almonds, Grapes, etc.)	At the beginning of blooming, fruit formation, fruit growing and before harvest.	250-300 cc in 100 liters of water	1000-1500 cc/da or 75-100 cc/tree
Bananas, Citrus, Figs, Olives, Walnuts, Almonds, Kiwis	At the beginning of blooming, fruit formation, fruit growing and before harvest.	250-300 cc in 100 liters of water	1000-1500 cc/da or 75-100 cc/tree
Industrial Crops and Legume (Sunflowers, Corns, Chickpeas, Soy, Beans, Peas, Peanuts, Lentil, Tea)	After the plants reach 10-15 cm in height, it is applied 2-3 times with an interval of 20 days.	200-250 cc in 100 liters of water	1000-1500 cc/da
Field Crops (Barley, Wheat, Rye, Paddy)	It is applied during tillering and bolting period.	200-250 cc in 100 liters of water	-----
Green Fields and Forage Plants (Clover, Sainfoin, Vetch)	It is applied 10 days after each mowing.	200-250 cc in 100 liters of water	-----

AGRIEGE TARIM GIDA İNŞ. İTH. İHR. SAN. ve TİC. LTD. ŞTİ.

ADRES: GAZI OSMAN PAŞA MAHALLESİ 5476 SOKAK NO:1

BUHARA İŞ MERKEZİ KAT:2 DAİRE:21 BORNOVA / İZMİR

TEL: 0232 483 61 22 • GSM: 0535 506 33 23

www.agriegie.com.tr • info@agriegie.com.tr

 www.linkedin.com/company/agriegie-tarim

 www.instagram.com/agriegieofficial

 www.facebook.com/agriegieofficial

 www.twitter.com/agriegieofficial