



## KENTON

Kenton is a liquid micro element containing 8% iron. Thanks to its special formula, it is effective for a long time in calcareous and alkaline soils. It prevents iron deficiency in vegetables, vineyards, citrus fruits, fruit trees and ornamental plants. Eliminates yellowing caused by iron deficiency.

GUARANTEED SUBSTANCE	% W/W
Water Soluble Iron (Fe)	8

### AREA, METHOD, TIME AND AMOUNT OF USE:

USAGE AREA	USAGE TIME	USAGE AND AMOUNT	
		BY LEAVES	BY SOIL
All Greenhouse Vegetables (Tomato, Pepper, Zucchini, Eggplant, Cucumber and Strawberry)	From planting, weekly application is carried out throughout the season.	100–150 cc for 100 liters of water	450–500 cc / da
All Open Field Vegetables (Tomato, Pepper, Zucchini, Eggplant, Cucumber and Strawberry)	After the seedlings are transplanted, the application is made every 15 days throughout the season.	100–150 cc for 100 liters of water	550–600 cc / da
All Winter Vegetables Curly, Lettuce, Leek, Spinach, Iceberg, Cabbage	Application is made throughout the season every 15 days from the seedling period.	10H–150 cc for 100 liters of water	550–600 cc / da
Cut Flowers	From planting, weekly practice is carried out throughout the season.	100–150 cc for 100 liters of water	450- 500 cc / da
All Fruit Trees (Apple, Pear, Cherry, Apricot, Quince)	At the beginning of flowering, during fruit drop and before fruit growing and harvest.	15H–200 cc for 100 liters of water	600-750 cc / da or 50 cc/ tree
Banana and Citrus, Fig, Olive, Walnut, Almond and Kiwi	At the beginning of flowering, during fruit drop and before fruit growing and harvest.	150–200 cc for 100 liters of water	600—750 cc / da or 50 cc/ tree
Viticulture,	It is applied after flowering, in the verdant period, before fruit growing period and harvest period.	150–200 cc for 100 liters of water	550—600 cc / da or 10 cc/ omca
Melon, Watermelon, Pumpkin etc.	It is applied throughout the season with an interval of 21 days after the second hoeing.	150 -200 cc for 100 liters of water	550-600 cc / da
All Tubular Plants (Onion, Potato, Turnip, Carrot, S. Beet, Garlic)	It is applied throughout the season with an interval of 21 days after the second hoeing.	15H-200 cc for 100 liters of water	550-600 cc / da
All Legumes (Chickpea, Lentil Beans, Soy Beans, Peanuts)	It is applied 2-3 times with 21 days intervals after the second hoeing.	150–200 cc for 100 liters of water	550—600 cc / da
All Industrial Plants (Cotton, Corn, Tea, Sunflower, Tobacco)	It is applied 2-3 times with 21 days intervals after the second hoeing.	150–200 cc for 100 liters of water	550-600 cc / da
All Field Crops (Wheat, Barley, Paddy, Oats)	It is applied with weed medicine and during tillering period.	150–200 cc for 100 liters of water	
All Green Field and Forage Plants (Clover, Sainfoin, Vetch)	It is applied ten days after each mowing.	150–200 cc for 100 liters of water	